

## Protocol for cryopreservation of HDMVEC/TERT164-B

page 1 of 2

Version: May 2021

Evercyte Ord. No.:	CHT-013-0164-B
Designation:	HDMVEC/TERT164-B, human dermal microvascular endothelial cells
Freezing medium:	HDMVEC/TERT164-B growth medium (see protocol <i>Passaging of HDMVEC/TERT164-B</i> ) 10 % FBS (Sigma Aldrich, Cat# F7524) 10 % DMSO (Sigma Aldrich, Cat# D2650)
<p>Preparation of 10 ml freezing medium, prepare just before use:</p> <ul style="list-style-type: none"><li>- take 8 ml of HDMVEC/TERT164 growth medium, transfer to 15 ml centrifugation tube</li><li>- add 1 ml of FBS</li><li>- add 1 ml of DMSO</li><li>- mix properly and store at 4°C</li></ul>	
Additional reagents:	0.1 % Gelatin (Sigma Aldrich, Cat# G1393, 2 %) 0.05 % Trypsin-EDTA (Gibco, Cat# 25300-054) Defined Trypsin Inhibitor (Gibco, Cat# R007100) PBS (Sigma Aldrich, Cat# D8537)
Freezing cells:	<ul style="list-style-type: none"><li>- detach the cells from the culture vessel by using Trypsin-EDTA solution as described in protocol <i>Passaging of HDMVEC/TERT164-B cells</i></li><li>- resuspend the detached cells in growth medium and centrifuge at 170 g for 5 min</li><li>- discard the supernatant</li><li>- resuspend the resulting cell pellet in the remaining droplet</li><li>- add freezing medium (tempered to 4°C) to reach a cell density of about <math>5 \times 10^5</math> cells/ml (for thawing in a 25 cm<sup>2</sup> culture flask)</li><li>- add 1 ml of this cell suspension to each pre-cooled cryovial and immediately transfer the cells to -80°C</li><li>- after 24 hours transfer the vials to the liquid nitrogen tank</li></ul>
Thawing cells:	<p><b>When you start cultivating the cells, please transfer the content of the original Evercyte vial containing HDMVEC/TERT164-B cells into a T25 roux flask as described in the following:</b></p> <ul style="list-style-type: none"><li>- pre-coat a 25 cm<sup>2</sup> culture flask with Gelatin solution as described in protocol <i>Passaging of HDMVEC/TERT164-B cells</i></li><li>- add 6 ml of growth medium to a 25 cm<sup>2</sup> culture flask and place the culture flask in the incubator for at least 30 min to allow the medium to reach 37°C and its normal pH</li><li>- take a vial of frozen cells, rinse it outside with ethanol and pre-warm in the hand until one last piece of frozen cells is seen</li><li>- then, immediately transfer the content of the vial to a 15 ml centrifugation tube filled with 9 ml of medium pre-cooled to 4°C and centrifuge for 5 min at 170 g</li></ul>

- discard the supernatant and resuspend the cell pellet in the remaining droplet
- add 1 ml of the pre-warmed medium to the cells, transfer the cells to the prepared culture flask and incubate at 37°C in a suitable incubator
- perform a medium change 24 hours after thawing, if the cells are already 80 % confluent at this point, they have to be passaged as described in protocol *Passaging of HDMVEC/TERT164-B cells*

## Related products:

HDMVEC/TERT164-B, microvascular endothelial cells (Evercyte, Cat# CHT-013-0164-B )  
HUVEC/TERT2, umbilical vein endothelial cells (Evercyte, Cat# CHT-006-0008)