

Product-Data-Sheet for HDF/TERT164

Version: May 2021

Evercyte Ord. No.:	CHT-008-0164
Designation:	HDF/TERT164
Biosafety Level:	1
Shipped:	Frozen on dry ice
Growth medium:	DMEM/Ham´s F12 supplemented with FBS and G418: <u>Final components:</u> DMEM/Ham´s F12 (1:1) (PAN Biotech, Cat# P04-41150) 10 % FBS (Sigma-Aldrich, Cat# F7524) 100 µg/ml G418 (InvivoGen, Cat# ant-gn-5)
Growth:	Adherent
Organism:	Homo sapiens (human)
Morphology:	Fibroblastoid, spindle shaped morphology
Source:	Adult skin (female)
Cell Type:	Fibroblasts
Antigen Expression:	Positive for vimentin, responds to TGF-β treatment by induction of Col1A1 expression
Ethical statement:	Approved by Institutional Review Board (IRB) in accordance with the Declaration of Helsinki
Comments:	<p>HDF/TERT164 was developed from human dermal fibroblasts transduced with a retroviral expression vector (pLXSN) containing the hTERT gene.</p> <p>The cells show the typical fibroblast morphology and express typical fibroblastoid markers.</p> <p>HDF/TERT164 cells can be grown for a minimum of 55 PDs without showing signs of growth retardation or replicative senescence. Cells readily recover from cryopreservation and no changes in growth characteristics have been observed after thawing.</p>
Propagation:	Cells are grown in DMEM/Ham´s F12 medium supplemented with FBS and G418 (see above) at 37°C in a humidified atmosphere with 5 % CO ₂ .
Subculturing:	For detachment of the cells remove and discard the culture medium and wash the cells once with PBS (160 µl/cm ²). Remove PBS completely. Then, add 0.05 % Trypsin-EDTA solution (room-temperature; 20 µl/cm ² ; Gibco, Cat# 25300-054), make sure that all cells have been in contact with this solution and incubate the culture flask at 37°C for

approximately 3 min. Observe cell detachment under an inverted microscope. As soon as all cells are detached (if necessary, agitate the cells by gently hitting the flask), add growth medium (about 160 $\mu\text{l}/\text{cm}^2$) and aspirate the cells by pipetting.

Add appropriate aliquots of the cell suspension to new culture vessels supplemented with growth medium (final volume of 240 $\mu\text{l}/\text{cm}^2$). A split ratio of 1:2 to 1:3 twice a week is recommended (after cells have reached about 90 % confluence).

Preservation:	<p>Freezing medium:</p> <p>DMEM/Ham's F12 (1:1) (PAN Biotech, Cat# P04-41150) 10 % FBS (Sigma-Aldrich, Cat# F7524) 10 % DMSO (Sigma-Aldrich, Cat# D2650)</p> <p>Storage temperature: liquid nitrogen</p>
Freezing and thawing procedure:	<p>Freezing of cells:</p> <p>Detach the cells from the culture vessel by using Trypsin-EDTA as described above, resuspend the detached cells in growth medium and centrifuge at 170 g for 5 min. Then, discard the supernatant, resuspend the resulting cell pellet in the remaining droplet and add freezing medium (tempered to 4°C) to reach a cell density of about 5×10^5 cells/ml (for thawing in a 25 cm^2 culture flask). Add 1 ml of this cell suspension to each pre-cooled cryovial and immediately transfer the cells to -80°C. After 24 hours transfer the vials to the liquid nitrogen tank.</p> <p>Thawing of cells:</p> <p>When you start cultivating the cells, please transfer the content of the original Evercyte vial containing HDF/TERT164 cells into a T25 roux flask as described in the following:</p> <p>Add 6 ml of growth medium to a 25 cm^2 culture flask and place the culture flask in the incubator for at least 30 min to allow the medium to reach 37°C and its normal pH. Take a vial of frozen cells, rinse it outside with ethanol and pre-warm in the hand until one last piece of frozen cells is seen. Then, immediately transfer the content of the vial to a 15 ml centrifugation tube pre-filled with 9 ml of medium pre-cooled to 4°C and centrifuge for 5 min at 170 g. Then, discard the supernatant and resuspend the cell pellet in the remaining droplet. Add 1 ml of the pre-warmed medium to the cells, transfer them to the prepared culture flask and incubate at 37°C in a suitable incubator.</p> <p>Perform a medium change 24 hours after thawing. If the cells are already 80-90 % confluent at this point, they have to be passaged (see above or protocol <i>Passaging of HDF/TERT164 cells</i>).</p>
Doubling Time:	60 - 72 hours
Virus Testing	Cells have been tested negative for HAV and Parvo B19 with Roche DPX-PCR (cobas® TaqScreen DPX-Test), for HBV, HCV, HIV nucleic acids with Roche-Multiplex-PCR (cobas® TaqScreen MPX Test, v2.0).

Other Analytical Data: Cells are negative for Mycoplasma contaminations as tested using MycoAlert™ Mycoplasma Detection Kit from Lonza.
Cells are negative for bacterial and fungal contaminations as tested according to Ph. Eur. 2.6.1. / USP <71>.
STR profile has been analyzed and is as expected.

Please Note:

The classification of biosafety level is based on Austrian Legislation (Gentechnikbuch; Systemverordnung) and on recommendations of the Central Committee on Biological Safety (ZKBS). While Evercyte undertakes all reasonable measures to test for absence of a selected panel of known human pathogenic viruses, there is currently no test procedure available that guarantees for complete absence of infectious pathogens. The use of state-of-the art infectious virus assays or viral antigen assays may leave open the possible existence of a latent viral genome, even if a negative test result is obtained. Therefore, we recommend that all human cell lines should be handled with caution such as an organism of ACDP Hazard Group 2. People who work with our cells must follow national regulations and safety precautions. The laboratories must be equipped with a security level according to the classification of the cells / products. Evercyte assumes no liability whatsoever in connection with the receipt, handling or the consequences of improper use of our products.

Notices and disclaimers: Evercyte products are intended for laboratory research purposes only. They are not intended for use in humans.

While Evercyte uses reasonable efforts to include accurate and up-to-date information on this site, no warranties or representations are made to its accuracy. Citations from scientific literature and patents are provided for informational purposes only and no warranties are made that such information has been confirmed to be accurate.