

## Product-Data-Sheet for HUVEC/TERT66

Version: May 2021

Evercyte Ord. No.:	CHT-006-0066
Designation:	HUVEC/TERT66
Biosafety Level:	1
Shipped:	Frozen on dry ice
Growth medium:	Endopan 300 SL Kit (PAN Biotech) supplemented with Panexin SL-S (PAN Biotech) and G418:  <u>Final components:</u> Endopan 300 SL (PAN Biotech, Cat# P04-0065K without GA) Serum substitute Panexin SL-S (PAN Biotech, Cat# P04-0065S) 20 µg/ml G418 (InvivoGen, Cat# ant-gn-5)
Growth:	Adherent
Organism:	Homo sapiens (human)
Morphology:	Endothelial
Source:	Umbilical cord
Cell Type:	Umbilical vein endothelial cells
Antigen Expression:	Positive for vWF, CD31
Ethical statement:	Approved by Institutional Review Board (IRB) in accordance with the Declaration of Helsinki
Comments:	HUVEC/TERT66 was developed from human umbilical vein endothelial cells by transduction with a retroviral expression vector (pLXSN) containing the hTERT gene. The cells can be grown under serum-free cell culture conditions. The cells form neoangiogenic webs on Matrigel.
Propagation:	Cells are grown in Endopan 300 SL medium supplemented with Panexin SL-S and G418 (see above) at 37°C in a humidified atmosphere with 5 % CO <sub>2</sub> .
Subculturing:	The new culture flasks have to be pre-coated with Gelatin (Sigma-Aldrich, Cat# G1393, diluted to 0.1 % with PBS). Therefore, the culture flasks are treated with 0.1 % Gelatin solution (80 µl/cm <sup>2</sup> ) at 37°C for at least 10 min (10-60 min). Before introducing cells, remove excess of Gelatin solution. For detachment of the cells remove and discard the culture medium and wash the cells twice with PBS (each 160 µl/cm <sup>2</sup> ). Remove PBS completely. Then, add 0.05 % Trypsin-EDTA solution (room-temperature; 20 µl/cm <sup>2</sup> ;

Gibco, Cat# 25300-054), make sure that all cells have been in contact with this solution and incubate the culture flask at 37°C for approximately 3 min. Observe cell detachment under an inverted microscope. As soon as all cells are detached (if necessary, agitate the cells by gently hitting the flask), add Defined Trypsin Inhibitor (20 µl/cm<sup>2</sup>). Thereafter, resuspend the cells in growth medium (about 160 µl/cm<sup>2</sup>), aspirate the cells by pipetting and centrifuge at 170 g for 5 min. Discard the supernatant, resuspend the cell pellet in the remaining droplet and add growth medium. Then, add appropriate aliquots of the cell suspension to Gelatin coated culture vessels supplemented with growth medium (final volume of 240 µl/cm<sup>2</sup>). A split ratio of 1:2 twice a week is recommended (after cells have reached about 90-95 % confluence). Cultivate cells at 37°C in a humidified atmosphere with 5 % CO<sub>2</sub>.

---

**Preservation:****Freezing medium:**

CryoStor<sup>®</sup> cell cryopreservation medium CS10 (Sigma-Aldrich, Cat# C2874)

Storage temperature: liquid nitrogen

---

**Freezing and thawing procedure:****Freezing of cells:**

Detach the cells (about 90-95 % confluence) from the culture vessel by using Trypsin-EDTA and Defined Trypsin Inhibitor as described above, resuspend the detached cells in growth medium and centrifuge at 170 g for 5 min. Then, discard the supernatant, resuspend the resulting cell pellet in the remaining droplet and add freezing medium (tempered to 4°C) to reach a cell density of about 5 x 10<sup>5</sup> cells/ml (for thawing in a 25 cm<sup>2</sup> culture flask). Add 1 ml of this cell suspension to each pre-cooled cryovial and immediately transfer the cells to -80°C. After 24 hours transfer the vials to the liquid nitrogen tank.

**Thawing of cells:**

**When you start cultivating the cells, please transfer the content of the original Evercyte vial containing HUVEC/TERT66 cells into a T25 roux flask as described in the following:**

Before thawing the original vial containing Evercyte cells, pre-coat a 25 cm<sup>2</sup> culture flask with Gelatin (see above and protocol *Passaging of HUVEC/TERT66 cells*). Add 6 ml of growth medium to a 25 cm<sup>2</sup> culture flask and place it in the incubator for at least 30 min to allow the medium to reach 37°C and its normal pH. Take a vial of frozen cells, rinse it outside with ethanol and pre-warm in the hand until one last piece of frozen cells is seen. Then, immediately transfer the content of the vial to a 15 ml centrifugation tube pre-filled with 9 ml of medium pre-cooled to 4°C and centrifuge for 5 min at 170 g. Then, discard the supernatant and resuspend the cell pellet in the remaining droplet. Add 1 ml of the pre-warmed medium to the cells, transfer them to the prepared culture flask and incubate at 37°C in a suitable incubator.

Perform a medium change 24 hours after thawing. If the cells are already 80 % confluent at this point, they have to be passaged (see above or protocol *Passaging of HUVEC/TERT66 cells*).

Doubling Time:	About 72 hours
Virus Testing	Cells have been tested negative for HAV and Parvo B19 with Roche DPX-PCR (cobas® TaqScreen DPX-Test), for HBV, HCV, HIV nucleic acids with Roche-Multiplex-PCR (cobas® TaqScreen MPX Test, v2.0).
Other Analytical Data:	Cells are negative for Mycoplasma contaminations as tested using MycoAlert™ Mycoplasma Detection Kit from Lonza. Cells are negative for bacterial and fungal contaminations as tested according to Ph. Eur. 2.6.1. / USP <71>. STR profile has been analyzed and is as expected.

**Please Note:**

The classification of biosafety level is based on Austrian Legislation (Gentechnikbuch; Systemverordnung) and on recommendations of the Central Committee on Biological Safety (ZKBS). While Evercyte undertakes all reasonable measures to test for absence of a selected panel of known human pathogenic viruses, there is currently no test procedure available that guarantees for complete absence of infectious pathogens. The use of state-of-the art infectious virus assays or viral antigen assays may leave open the possible existence of a latent viral genome, even if a negative test result is obtained. Therefore, we recommend that all human cell lines should be handled with caution such as an organism of ACDP Hazard Group 2. People who work with our cells must follow national regulations and safety precautions. The laboratories must be equipped with a security level according to the classification of the cells / products. Evercyte assumes no liability whatsoever in connection with the receipt, handling or the consequences of improper use of our products.

Notices and disclaimers: Evercyte products are intended for laboratory research purposes only. They are not intended for use in humans.

While Evercyte uses reasonable efforts to include accurate and up-to-date information on this site, no warranties or representations are made to its accuracy. Citations from scientific literature and patents are provided for informational purposes only and no warranties are made that such information has been confirmed to be accurate.