

Product-Data-Sheet for ASC/TERT1

Version: March 2023

Evercyte Ord. No.:	CHT-001-0005
Designation:	ASC/TERT1
Biosafety Level:	1
Shipped:	Frozen on dry ice
Growth medium:	EGM TM -2 Endothelial Cell Growth Medium (Lonza, Cat# CC-3162) supplemented with FBS and G418: <u>Final components:</u> EBM TM -2 basal medium (Lonza, Cat# CC-3156) Components of EGM TM -2 SingleQuots TM (Lonza, Cat# CC-4176: Hydrocortisone, hEGF, VEGF, R3-IGF-1, Ascorbic Acid, hEGF, Heparin) 4 % FBS (PAN Biotech, Cat# P30-3031, ready-to-use, stored at 4°C after thawing) 200 µg/ml G418 (InvivoGen, Cat# ant-gn5, ready-to-use, stored at -20°C)
Growth:	Adherent
Organism:	Homo sapiens (human)
Morphology:	Mesenchymal
Source:	Adipose tissue / liposuction (female donor)
Cell Type:	Mesenchymal stromal cells
Antigen Expression:	Positive for CD73, CD90, CD105; negative for CD34
Ethical statement:	Approved by Institutional Review Board (IRB) in accordance with the Declaration of Helsinki
Comments:	<p>ASC/TERT1 was developed from human adipose tissue-derived mesenchymal stromal cells by transduction with a retroviral expression vector (pLXSN) containing the hTERT gene.</p> <p>Cells show expression of CD73, CD90 and CD105 and are negative for CD34. Cells can be differentiated towards adipogenic, osteogenic and chondrogenic lineages and are characterized by immunomodulatory properties.</p> <p>The cell line was continuously cultured for more than 70 population doublings without showing signs of growth retardation or replicative senescence whereas the parental cells entered senescence after having reached around 10 population doublings.</p>
Propagation:	Cells are grown in EGM TM -2 Endothelial Cell Growth Medium supplemented with FBS and G418 (see above) at 37°C in a humidified atmosphere with 5 % CO ₂ .

Subculturing:	<p>For detachment of the cells remove and discard the culture medium and wash the cells twice with PBS. Remove PBS completely. Then, add 0.05 % Trypsin-EDTA solution (RT, 20 $\mu\text{l}/\text{cm}^2$, Gibco, Cat# 25300054), make sure that all cells have been in contact with this solution and incubate the culture flask at 37°C for approximately 2-3 min. Observe cell detachment under an inverted microscope. As soon as all cells are detached (if necessary, agitate the cells by gently hitting the flask), add Defined Trypsin Inhibitor (20 $\mu\text{l}/\text{cm}^2$, Gibco, Cat# R007100).</p> <p>Thereafter, resuspend the cells in growth medium (about 160 $\mu\text{l}/\text{cm}^2$) and aspirate the cells by pipetting, centrifuge at 170 g for 5 min. Discard the supernatant, resuspend the cell pellet in the remaining droplet and add growth medium. Then, transfer appropriate aliquots of the cell to new culture vessels supplemented with growth medium (final volume of 240 $\mu\text{l}/\text{cm}^2$). A split ratio of 1:3 to 1:4 twice a week is recommended (after having reached about 80 % confluence).</p>
Preservation:	<p>Freezing medium:</p> <p>ASC/TERT1 growth medium (see above) 10 % FBS (Sigma-Aldrich, Cat# F7524, ready-to-use, stored at 4°C after thawing) 10 % DMSO (Sigma-Aldrich, Cat# D2650, ready-to-use, stored at RT)</p> <p>Storage temperature: liquid nitrogen</p>
Freezing and thawing procedure:	<p>Freezing of cells:</p> <p>Detach the cells from the culture vessel by using Trypsin-EDTA and Defined Trypsin Inhibitor as described above, resuspend the detached cells in growth medium and centrifuge at 170 g for 5 min. Then, discard the supernatant, resuspend the resulting cell pellet in the remaining droplet and add freezing medium (tempered to 4°C) to reach a cell density of about 5×10^5 cells/ml (for thawing in a 25 cm^2 culture flask). Add 1 ml of this cell suspension to each pre-cooled cryovial and immediately transfer the cells to -80°C. After 24 hours transfer the vials to the liquid nitrogen tank.</p> <p>Thawing of cells:</p> <p>When you start cultivating the cells, please transfer the content of the original Evercyte vial containing ASC/TERT1 cells into a T25 roux flask as described in the following:</p> <p>Add 6 ml of growth medium to a 25 cm^2 culture flask and place the culture flask in the incubator for at least 30 min to allow the medium to reach 37°C and its normal pH. Take a vial of frozen cells, rinse it outside with ethanol and pre-warm in the hand until one last piece of frozen cells is seen. Then, immediately transfer the content of the vial to a 15 ml centrifugation tube pre-filled with 9 ml of growth medium pre-cooled to 4°C and centrifuge for 5 min at 170 g. Then, discard the supernatant and resuspend the cell pellet in the remaining droplet. Add 1 ml of the pre-warmed medium to the cells, transfer them to the prepared culture flask and incubate at 37°C in a suitable incubator.</p>

Perform a medium change 24 hours after thawing. If the cells are already 80 % confluent at this point, they have to be passaged (see above or protocol *Passaging of ASC/TERT1 cells*).

Doubling Time:	36-48 hours
Virus Testing	Cells have been tested negative for HAV and Parvo B19 with Roche DPX-PCR (cobas® TaqScreen DPX-Test), for HBV, HCV, HIV nucleic acids with Roche-Multiplex-PCR (cobas® TaqScreen MPX Test, v2.0).
Other Analytical Data:	Cells are negative for Mycoplasma contaminations as tested using MycoAlert™ Mycoplasma Detection Kit from Lonza. Cells are negative for bacterial and fungal contaminations as tested according to Ph. Eur. 2.6.1. / USP <71>. STR profile has been analyzed and is as expected.

Please Note:

The classification of biosafety level is based on Austrian Legislation (Gentechnikbuch; Systemverordnung) and on recommendations of the Central Committee on Biological Safety (ZKBS). While Evercyte undertakes all reasonable measures to test for absence of a selected panel of known human pathogenic viruses, there is currently no test procedure available that guarantees for complete absence of infectious pathogens. The use of state-of-the art infectious virus assays or viral antigen assays may leave open the possible existence of a latent viral genome, even if a negative test result is obtained. Therefore, we recommend that all human cell lines should be handled with caution such as an organism of ACDP Hazard Group 2. People who work with our cells must follow national regulations and safety precautions. The laboratories must be equipped with a security level according to the classification of the cells / products. Evercyte assumes no liability whatsoever in connection with the receipt, handling or the consequences of improper use of our products.

Notices and disclaimers: Evercyte products are intended for laboratory research purposes only. They are not intended for use in humans.

While Evercyte uses reasonable efforts to include accurate and up-to-date information on this site, no warranties or representations are made to its accuracy. Citations from scientific literature and patents are provided for informational purposes only and no warranties are made that such information has been confirmed to be accurate.